



# Reducing the conflict between Cormorants and fisheries on a pan-European scale

REDCAFE

Final Report

Report of a Concerted Action funded by the European Union. Study contract no. Q5CA-2000-31387: Reducing the conflict between cormorants and fisheries on a pan-European scale.

Edited by D N Carss

Centre for Ecology & Hydrology Banchory, Hill of Brathens, Banchory  
Aberdeenshire, AB31 4BW, Scotland, UK.

# Natural Environment Research Council Centre for Ecology & Hydrology

**CEH Banchory**  
Hill of Brathens  
Banchory  
Aberdeenshire  
AB31 4BY  
Scotland, UK.

Fax: +44 1330 823303  
Internet: d.carss@ceh.ac.uk  
Web: www.banchory.ceh.ac.uk

CEH Contract Number: C01749

## Reducing the conflict between cormorants and fisheries on a pan-European scale Final Report

Edited by: D N Carss

Commissioned by: European Commission  
DG XIV Directorate-General for Fisheries  
Rue de la Loi 200  
Bâtiment J II 99 6/11  
B-1049 Brussels, Belgium

Contract number: Q5CA-2000-31387

*This report does not necessarily reflect the views of the European Commission and in no way anticipates any future opinion of the Commission. The contents of the report may not be reproduced unless the source of the material is indicated.*

*This study has been carried out with the financial assistance of the European Commission.*

This report is drafted at the request of the commissioner indicated above and is his property. Nothing from this report may be reproduced and/or published by print, photoprint microfilm or any other means without the previous written consent from the commissioner of the study.

Photographs on the cover (Top left, clockwise): Great cormorant © David Grémillet, coastal fisheries, Greece © Dave Carss, recreational fisherman © Trout & Salmon magazine, Danish pound net fishermen, © DC, Roach © Roger Phillips and Martyn Rix, REDCAFE participants, Waltham Abbey, November 2002, © Szymon Bzoma.

## PARTICIPATING ORGANISATIONS

<p>David Carss (co-ordinator, steering group)  Centre for Ecology &amp; Hydrology  CEH Banchory, Hill of Brathens,  Banchory, Aberdeenshire, AB31 4BW  United Kingdom.  d.carss@ceh.ac.uk</p>
<p>Ian Winfield  Centre for Ecology &amp; Hydrology  CEH Windermere, Far Sawrey,  Ambleside, Cumbria, LA22 0LP  United Kingdom.  i.winfield@ceh.ac.uk</p>
<p>Phillip Scott Jones  University of Wolverhampton  Centre for Rural Development and Training,  WS1 3BD, United Kingdom.  p.s.jones@wlv.ac.uk</p>
<p>Mariella Marzano  CEH Banchory (<a href="mailto:mama@ceh.ac.uk">mama@ceh.ac.uk</a>) and  Department of Anthropology  University of Durham  43 Old Elvet  Durham, DH1 3HN  United Kingdom.  mariella.marzano@durham.ac.uk</p>
<p>Morten Frederiksen  CEH Banchory, Hill of Brathens,  Banchory, Aberdeenshire, AB31 4BW  United Kingdom  mfr@ceh.ac.uk</p>
<p>Erik Petersson  National Board of Fisheries  Institute of Freshwater Research,  Staangholmsvaegen, S-178 93,  Drottningholm,  Sweden.  erik.petersson@fiskeriverket.se</p>
<p>Henri Engström  Ex University of Uppsala, Dept. of Population Biology  Now: SKB, Swedish Nuclear Fuel and Waste Management Co  Platsundersökning i Forsmark  742 03 Östhammar, Sweden.  henri.engstrom@skb.se</p>
<p>Thomas Bregnballe (steering group)  National Environmental Research Institute  Dept. of Coastal Zone Ecology. Kaloe, Genaavej 12,  DK-8410, Roende, Denmark.  tb@dmu.dk</p>
<p>Thomas Keller (steering group)  The Technical University of Munich  Technische Universitaet Muenchen,  Institut fuer Tierwissenschaften, Angewandte Zoologie,  Alte Akademie 16, D-85350, Freising  Germany.  t.keller@lrz.tum.de</p>

<p>Mennobart van Eerden (steering group)  Institute for Inland Water Management and Waste Water Treatment  RIZA, Directoraat-General Jijkswaterstaat, PO Box 17, NL-8200,  Lelystad, The Netherlands.  <a href="mailto:m.vEerden@riza.rws.minvenw.nl">m.vEerden@riza.rws.minvenw.nl</a></p>
<p>Loïc Marion  Department of Ecology, University of Rennes  Université de Rennes 1, ECOBIO-UMR 6553,  U.F.R. S.V.E., CS74205, Avenue du Général Leclerc 236,  F-35042, Rennes, France.  <a href="mailto:loic.marion@univ-rennes1.fr">loic.marion@univ-rennes1.fr</a></p>
<p>Daniel Gerdeaux  Institute of Lacustrine Hydrobiology  Institut National de la Recherche Agronomique,  Station d'Hydrobiologie Lacustre, BP R11,  75 Avenue de Corzent, F-74203, Thonon-Les-Bains,  France.  <a href="mailto:gerdeaux@thonon.inra.fr">gerdeaux@thonon.inra.fr</a></p>
<p>David Grémillet  Centre for Ecological and Physiological Energetics  Centre d'Ecologie et Physiologie Energetiques,  23 Rue Becquerel, F-67087, Strasbourg,  France.  <a href="mailto:david.gremillet@c-strasbourg.fr">david.gremillet@c-strasbourg.fr</a></p>
<p>Carlos Garcia de Leaniz, Angel Serdio &amp; Sofía Consuegra  Direccion General de Montes y Conservacion de la Naturaleza  Institute of Freshwater Fisheries - Arredondo,  Arturo Lopez s/n, E-39813, Arredondo, Cantabria,  Spain.  <a href="mailto:centro_ictiologico_arredondo@yahoo.com">centro_ictiologico_arredondo@yahoo.com</a></p>
<p>Bruno Broughton  European Anglers Alliance  Trenchard, Lower Bromstead Road,  Moreton, Newport, Shropshire,  TF10 9DQ  United Kingdom.  <a href="mailto:bruno.broughton@virgin.net">bruno.broughton@virgin.net</a></p>
<p>Savas Kazantzidis (Maria Panayotopoulou, 2001)  Hellenic Ornithological Society  G. Rodiou 10 St., GR-546 38 Thessaloniki, Greece.  <a href="mailto:hrysavgi@otenet.gr">hrysavgi@otenet.gr</a></p>
<p>Erich Staub  European Inland Fisheries Advisory Commission  EIFAC, CH-3003, Bern, Switzerland.  <a href="mailto:erich.staub@buwal.admin.ch">erich.staub@buwal.admin.ch</a></p>
<p>Ido Izhaki &amp; Gadi Katzir  Department of Biology, University of Haifa  University of Haifa at Oranim, IL-36006, Tivon, Israel.  <a href="mailto:izhaki@research.haifa.ac.il">izhaki@research.haifa.ac.il</a></p>
<p>Zeef Arad  Department of Biology, Institute of Technology  Technion – Israel Institute of Technology – Faculty of Biology,  Technion City, IL-32000, Israel.  <a href="mailto:zarad@techunix.technion.ac.il">zarad@techunix.technion.ac.il</a></p>
<p>Tamir Strod &amp; Jonathan Harari  The Hula Nature Reserve  P.O.B. 340 Yesud Hamaala,  12105, Israel.  <a href="mailto:st.hula@nature-parks.org.il">st.hula@nature-parks.org.il</a></p>

<p>Harald Kleisinger  Bavarian Bureau of Environmental Protection  Lfu Bayern, Nature Protection and Landscape Planning,  Buergermeister-Ulrich-Strasse 160, D-86179, Augsburg,  Germany.  <a href="mailto:harald.kleisinger@lfu.bayern.de">harald.kleisinger@lfu.bayern.de</a></p>
<p>Julian Hughes  Royal Society for the Protection of Birds  RSPB, Species Policy Unit,  The Lodge, Sandy, Bedfordshire, SG19 2DL,  United Kingdom.  <a href="mailto:julian.hughes@rspb.org.uk">julian.hughes@rspb.org.uk</a></p>
<p>Ian Russell  The Centre for Environment, Fisheries &amp; Aquaculture Science  CEFAS Lowestoft Laboratory, Pakefield Road,  Lowestoft, Suffolk, NR33 0HT,  United Kingdom.  <a href="mailto:i.c.russell@cefas.co.uk">i.c.russell@cefas.co.uk</a></p>
<p>Robert Gwiazda  Institute of Freshwater Biology, Polish Academy of Sciences  Polish Academy of Sciences, Slawkowska 17, Krakow,  PL-31-016,  Poland.  <a href="mailto:gwiazda@zbw.pan.krakow.pl">gwiazda@zbw.pan.krakow.pl</a></p>
<p>Redik Eschbaum  Estonian Marine Institute  Estonian Marine Institute, Viljandi Road 18B,  11216 Tallinn, Estonia.  <a href="mailto:eschbaum@ut.ee">eschbaum@ut.ee</a></p>
<p>Willem Dekker  Netherlands Institute for Fisheries Research RIVO-DLO  PO Box 68, NL-1970 AB IJmuiden,  The Netherlands.  <a href="mailto:Willem@rivo.wag-ur.nl">Willem@rivo.wag-ur.nl</a></p>
<p>Christian Dieperink  Ex. Danish Institute for Fisheries Research  Now: Waterframe, Hejnaesvej 4,  DK – 8680 Ry,  Denmark.  <a href="mailto:cd@waterframe.dk">cd@waterframe.dk</a></p>
<p>Janis Baumanis  Institute of Biology  Laboratory of Ornithology, Miera st. 3, LV-2169,  Salaspils, Latvia.  <a href="mailto:jbaumanis@email.lubi.edu.lv">jbaumanis@email.lubi.edu.lv</a></p>
<p>Nis Røv &amp; Svein Lorentsen  Norwegian Institute for Nature Research  Tungasletta 2, N-7485 Trondheim, Norway.  <a href="mailto:nils.rov@ninatrd.ninaniku.no">nils.rov@ninatrd.ninaniku.no</a>  <a href="mailto:shl@ninatrd.ninaniku.no">shl@ninatrd.ninaniku.no</a></p>
<p>Timo Asanti  Finnish Environmental Institute  Nature and land use Division, Kesäkuu 6, PO Box 140,  FIN-00251, Helsinki, Finland.  <a href="mailto:timo.asanti@vyh.fi">timo.asanti@vyh.fi</a></p>

<p>Petr Musil &amp; Renata Martincova  Charles University, Prague  Department of Zoology, Faculty of Sciences,  Vini ná 7, Praha 2, CV-128 44, Prague,  Czech Republic.  p.musil@post.cz  mar_ren@hotmail.com</p>
<p>Josef Trauttmansdorf  Otto Koenig Institute  A-2000, Stockerau, Austria.  <a href="mailto:okido@aon.at">okido@aon.at</a></p>
<p>Rosemarie Parz-Gollner  Inst. fuer Wildbiologie und Jagdwirtschaft  Univ. f. Bodenkultur Wien, Peter Jordanstrasse 76,  A-1190, Vienna, Austria.  Rosemarie.parz-gollner@boku.ac.at</p>
<p>Stefano Volponi  Ex University of Ferrara  Now: Istituto Nazionale Fauna Selvatica  Via Ca' Fornacetta, 9  40064 Ozzano Emilia BO  Italy.  svolponi@racine.ra.it</p>
<p>Jean-Yves Paquet  Central Ornithologique Aves  Rue de Blacet 1, 5530 Yvoir, Belgium.  <a href="mailto:jeanyves.paquet8@yucum.be">jeanyves.paquet8@yucum.be</a></p>
<p>Ger Rogan  Marine Institute  Salmon Management Services, Furnace,  Newport, Co. Galway, Ireland.  ger.rogan@marine.ie</p>
<p>Michael Andersen  Danish Fishermen's Association  H.C. Andersens Boulevard 37, PO Box 403,  DK-1504, Copenhagen, Denmark.  ma@fiskeriforening.dk</p>
<p>Szymon Bzoma  University of Gdańsk  Dept. of Vertebrate Ecology and Zoology,  Al. Legionów 9, 80-441, Gdańsk, Poland.  bioszb@univ.gda.pl</p>
<p>Linas Ložys  Laboratory of Marine Ecology  Institute of Ecology, Akademijos 2, LT-2600,  Vilnius, Lithuania.  lozys@eko.lt</p>
<p>Ivailo Nikolov  Ex Sofia University  Now: "Mladost" 4 bl 403, ent.1,  AP4, 1715 Sofia,  Bulgaria.  ivailo_nikolov@abv.bg</p>
<p>Vilju Lilleleht  Estonian Agricultural University  Institute of Zoology and Botany,  Riia 181, EE51014 Tartu, Estonia.  vilju.lilleleht@zbi.ee</p>

<p>Nikolay Kissiov  Bulgarian Fisheries and Aquaculture Association  j.k. "Levski-g" bd. 10-7-132, 1 836 Sofia, Bulgaria.  n.kissiov@asico.net</p>
<p>Botond Kiss  Danube Delta Institute  Str Babadag, Ro8800, Tulcea, Romania.  <a href="mailto:jbkiss@indd.tim.ro">jbkiss@indd.tim.ro</a></p>
<p>Miha Janc  Fisheries Association of Slovenia  Trzaska cesta 132,  Slovenia.  miha.janc@siol.net</p>
<p>Catarina Vinagre &amp; Susana Franca  University of Lisbon  Departamento de Zoologie e Anthropologia / Instituto de Oceanografia,  Faculdade de ciencias da Universidade de Lisboa  R. Ernesto de Vasconcelos, 1749-016 Lisbon  Portugal.  catv@fc.ul.pt  sfranca@fc.ul.pt</p>
<p>Kareen Seiche  On behalf of:  Saxon Ministry of Environment  Archivstrasse 1  01097 Dresden  Germany  kareen.seiche@t-online.de</p>

## 0 Executive summary

### 0.1 *Background to the project*

Two subspecies of Great Cormorant (hereafter ‘Cormorant’) occur in Europe: the ‘Atlantic’ subspecies *Phalacrocorax carbo carbo* and the ‘Continental subspecies *P. c. sinensis*. Latest (1995) breeding estimates for *carbo* are of 40,000 pairs, mostly on the coasts of Norway, UK, Ireland and northern France. The *sinensis* population (1995) is estimated to be over 150,000 pairs throughout the region, a dramatic increase since the 1960s. It is likely that the species is now more numerous across Europe than ever before. The geographical range of these populations has also expanded with Cormorants returning to some areas after a long absence and also moving into previously unoccupied area. The reasons for such expansion are unclear but possible causal factors include a “non-limiting food supply” and protective legislation, particularly EEC Directive 79/409 on the Conservation of Wild Birds. Cormorants are generalist fish-eating predators taking a wide variety of species in shallow coastal seas, running and standing freshwaters, and both traditional/extensive and intensive aquaculture systems. In almost all countries where Cormorants occur, their increasing numbers and geographical spread has led to a growing number of conflicts with commercial fisheries and recreational angling interests.

### 0.2 *Aims and set up of the project*

Although there are several national and/or international Cormorant management plans aimed at reducing such conflicts with Cormorants, there is no co-ordinated implementation at the international level and, in practice, and certainly for many affected by the ‘Cormorant problem’, these plans appear ineffectual. The REDCAFE project (December 2000 – November 2001) was designed to complement and develop previous work through synthesising available information on Cormorant conflicts and aspects of Cormorant ecology leading to them, through identifying methods of reducing the current Europe-wide conflict between Cormorants and fisheries interests and collating expert evaluations of their practical use. The project also addressed a specific Cormorant-fisheries conflict case study involving recreational angling in S. E. England. REDCAFE took a novel approach to delivering solutions to these problems by, for the first time, bringing together avian, fisheries and social scientists and many other relevant ‘stakeholders’ to discuss and report on these issues in a rigorous, co-ordinated and equitable manner. With these aims in mind, a pan-European network of project participants was established comprising 49 people representing 43 organisations from 25 countries and including seven main stakeholder groups: commercial fishermen, recreational fishermen, aquaculturists, avian/wetland conservationists, fisheries scientists, avian ecologists and social scientists.

### 0.3 *Cormorant conflicts with fisheries*

Various stakeholder groups often hold different values and, consequently, have different preferences for the use of limited natural resources: conflict is thus often inevitable. In addition to addressing environmental conflicts from a biological perspective, the social and cultural dimensions of human society that influence such conflicts also demand equal attention. Successful conflict management depends on conflicting parties opening communication channels and developing networks of trust for effective participation, dialogue and collaboration. Thus, wherever possible, information for the synthesis of Cormorant conflicts was provided by stakeholders affected directly by Cormorants. The provision and collation of information for the

present conflict synthesis formed the basis for REDCAFE's pan-European dialogue with stakeholders. This process also highlighted the difficulties involved in creating and managing dialogue between stakeholders from many countries and diverse backgrounds and these issues are discussed.

#### *0.4 Cases of Cormorant conflicts*

REDCAFE sampled Cormorant conflicts in 24 countries and collated information on 235 conflict cases. Cormorant conflicts were reported from a wide variety of habitats and fishery types: rivers, lakes, freshwater aquaculture ponds, coasts, and coastal aquaculture sites. This demonstrated the widespread geographical distribution of conflicts. Conflicts were reported by four different stakeholder groups representing recreational, commercial and nature conservation interests and covered a wide variety of fishery types, suggesting that the nature of conflicts also differed on a geographic scale.

#### *0.5 Habitat features of conflict cases*

Two species of cormorant were recorded in conflicts: both races of the Great Cormorant and the Pygmy Cormorant (*P. Pygmeus*). The geographical distributions of both species, as recorded in conflicts, followed closely their known breeding and/or wintering distributions. Cormorant conflicts were reported mostly from lower altitudes (< 500m). Within river systems, Cormorant conflicts on a pan-European scale showed similar distribution patterns. They were very much restricted to the lower and middle reaches, and hence relatively wide (i.e. 10-50m) stretches, of rivers. Similar, restricted distribution patterns were clear for conflict cases on the coast which were restricted to those localities with access to shallow (< 50m deep) inshore coastal water. Overall, most conflict cases were reported on nutrient-rich (i.e. eutrophic) waters, particularly freshwater aquaculture ponds, lakes and coasts, supporting the idea that Cormorant distribution is, in part at least, determined by the nutrient status of these waters.

#### *0.6 Conflicts in time and space*

Information on the seasonality of Cormorant conflicts showed patterns that fitted closely with the known seasonal movements of birds across Europe. As a consequence, the broad pan-European picture of Cormorant conflicts has three elements. First, winter (October-March) conflicts in those countries where birds overwinter, either towards the north west or south east. Second, summer (April-September) conflicts, presumably involving breeding birds, in the Netherlands and almost all countries bounding the Baltic. Third, conflicts throughout the year in the 'centre' of Europe (Denmark, Germany and the Czech Republic), presumably involving both breeding birds and others overwintering there from the north. Cormorant abundance increased with water surface area on a pan-European scale for stillwater lakes, freshwater aquaculture ponds and coasts and water surface area explained 56% of the variation in maximum Cormorant numbers across these habitats. There was no such relationship on rivers based on the information available for this synthesis. Such apparent differences require further investigation, particularly as information suggests that average Cormorant density on rivers is significantly higher than that in other habitats.

### 0.7 *Conflicts: fish*

Throughout Europe, there were strong associations between particular fish groups reported in conflict cases and particular habitat and fishery types. A wide variety of fish species were reported in relation to coastal conflicts. Cyprinids and salmonids were the main groups of fish recorded by stakeholders in relation to Cormorant conflicts on rivers. Similarly cyprinids, especially Carp, plus some salmonids, Perch and Pike were involved in conflicts at freshwater aquaculture ponds. Many conflicts were reported at Carp ponds throughout Europe and these sites are considered highly attractive to Cormorants in places such as the Czech Republic, Bavaria, southern Germany, and France. A small group of fishes including mullets, sea basses and sea breams were involved in conflicts at coastal, often extensive lagoon, aquaculture sites of southern Europe.

### 0.8 *Conflicts: finance*

Financial information was provided by fishery-related stakeholders for 105 conflict cases, approximately 45% of those recorded in the present synthesis. Nature conservation stakeholders did not provide any financial information in relation to any of the conflict cases they recorded. Fishery stakeholders provided information on the annual financial turnover in their fishery system and the turnover loss due to Cormorants as 'actual' figures or as 'estimates'(derived by unknown means), thus care must be taken when interpreting the financial information collected in this synthesis. Nevertheless, the 105 conflict cases gave a cumulative total for annual turnover of about 154 million euro and associated losses to Cormorants were given at about 17 million euro, an overall loss of 11%. There were significant differences in the scale of financial losses reported by the relevant stakeholders for different habitats and fishery types. All three fishery stakeholder groups independently were consistent in their views on relatively low financial losses due to Cormorants, recording average values of 9-12% of annual turnover. Around 2% of aquaculturist, 13% of commercial freshwater fishermen and 31% of commercial coastal fishermen recorded losses greater than 50% of the annual financial turnover in their fishery. In contrast, recreational anglers recorded considerably higher financial losses due to Cormorants, averaging 57% of annual turnover. Furthermore, in 43% of cases, anglers recorded financial losses greater than 50% of the annual turnover in their fishery. Although the disparity between commercial and recreational stakeholders' perceptions of financial losses due to Cormorants was clear from the information provided, the explanation for it was not and requires further investigation.

### 0.9 *Conflict issues*

Nine specific conflict issues were most commonly cited as being major ones for stakeholders. For both aquaculturists and commercial fishermen, the issue of **reduced catches** was most important whilst for both recreational anglers and nature conservationists the most important issue was **reduced fish stock through lowered production**. Recreational stakeholders also most frequently reported conflicts over reduced catches and **effects on fish population dynamics and community structure**, an issue that was also important to nature conservationists. Both aquaculturists and commercial fishermen were concerned over **loss of earnings from the fishery**, the former stakeholders cited conflicts over **loss of stocked fish** and the latter ones cited conflicts over reduced stock through lowered production. Finally, nature conservationists also frequently recorded concerns over **loss of juvenile fish and lowered recruitment, scaring/shooting disturbance, drowning of Cormorants**

**in fishing gear and damage to vegetation and landscape.** Thus, although stakeholder groups frequently shared concerns over specific major conflict issues, some concerns were specific to particular groups. Most importantly, nature conservationists cited broader ‘environmental’ issues more frequently than did the three fishery-related stakeholder groups. The conflict synthesis showed considerable, and consistent, similarities between the opinions of both income-producing stakeholder groups involved in fisheries. Although recreational anglers shared many of the concerns of these other fishery-related stakeholder groups, they also recorded some different major conflict issues. However, the biggest differences were between fishery-related stakeholders and nature conservationists. Nature conservationists, in general, were most concerned with wider (i.e. ‘environmental’) conflict issues.

#### *0.10 Information sources*

Stakeholders provided over 3, 500 records of the type of information they used to inform themselves about Cormorant conflict issues. Although most records were categorised as ‘popular’, this category included a range of diverse sources. Overall, only 15% of information sources used by stakeholders were assigned to the scientific literature. For all stakeholder groups, scientific literature was the least frequently recorded information source. The importance of ‘popular’ sources of information to all four stakeholder groups contributing to this synthesis was thus clear. For several specific conflict issues, different stakeholder groups claimed to be informed by scientific literature yet considered the magnitude of such conflicts to be very different. It is clear that there is a need for better dissemination of scientific information and for better understanding of the limitations and implications of scientific research.

#### *0.11 Cormorant ecology: factors leading to conflicts*

Any successful resolution, or management, of the conflicts between Cormorants and fisheries interests on a pan-European scale must include careful consideration of the best available biological information on Cormorant populations throughout the region. REDCAFE thus synthesised aspects of Cormorant ecology that lead conflicts. Relevant factors were categorised into four main themes: (1) general ecology and habitat features, (2) migration and the annual cycle, (3) fish communities and Cormorant diet, and (4) Cormorant ecology and impact at fisheries.

#### *0.12 Ecology synthesis in relation to Cormorants*

Cormorant ecology has been well studied. With respect to numbers, distribution, migratory movements, foraging behaviour and diet it is one of the best known wild bird in Europe. It is clear that Cormorants are opportunistic generalist fish predators. As a result of their broad ecological requirements, they do have the potential for considerable conflicts at specific fisheries. This is because, as well as flexibility in feeding site choice, generalist predators like the Cormorant could have considerable impact on their preferred prey species because their numbers are buffered to some extent against declines in these prey by their ability to switch to other types. The opportunistic nature of its foraging behaviour and its great adaptability to a variety of habitats, both freshwater and marine, makes the Cormorant an exceptionally successful species which is currently probably more abundant in western Europe than ever before and still expanding numerically in eastern Europe. This expansion in numbers and area is the result of European wide protective measures, eutrophication, the reduction of pesticides in the environment and alterations of water systems such as dams, sluices which facilitate foraging.

### *0.13 Ecology synthesis in relation to fish*

Fish species eaten by Cormorants are, for the most part, common, widespread species. The heavy fishery pressure exerted by people in many water systems in Europe has resulted in a shift in size distribution towards the smaller classes, which enhances Cormorant foraging conditions. Fewer large predatory fish are now present in many European waters because of over-fishing. This enables populations of smaller fish species to increase, which in turn favours the Cormorant. Eutrophication of water bodies has altered fish community - (and size -) structure again increasing the possibilities for Cormorants to exploit larger densities of small prey fishes.

### *0.14 Ecology synthesis in relation to damage at fisheries*

Fish species eaten by Cormorants are, for the most part, common, widespread species. The heavy fishery pressure exerted in many water systems in Europe has resulted in a shift in size distribution towards the smaller classes, which enhances Cormorant foraging conditions. Reduction of eutrophication will decrease Cormorant numbers through reduction in the carrying capacity of fishing waters. Restoration of waterways, aiming at a greater connectivity, will favour fish populations and reduce predation risk. In fish farming areas, specific knowledge on prey detection underwater may help to reduce predation of small fish. Enlarging stocked fish above the range commonly eaten by Cormorants (i.e. >500 g) may act to reduce the damage caused by birds. Periods of large-scale Cormorant movements through Europe (e.g. March and October) require extra management attention to avoid the establishment of any tradition to visit stocked water bodies or fish farm areas. A combination of ecological, demographic, climatological and geographical data into a GIS based Decision Support System may help to predict future Cormorant 'problems' and reduce current ones through integrated management.

### *0.15 Potential Cormorant management tools*

Potential Cormorant management tools were assessed on two spatial/temporal scales: long-term control of European Cormorants at the population level and shorter-term site-specific control measures. The synthesis aimed to provide a comprehensive overview of potential Cormorant management tools. It provides a review of population modelling and a synthesis of site-specific techniques and actions used against Cormorants. The synthesis also includes semi-quantitative information on the 'usefulness' of techniques in relation to their effectiveness (i.e. how long a technique works for), practicability (i.e. how easy the technique is to use), acceptability (i.e. how the technique is viewed by both stakeholders and the general public) and costs. REDCAFE participants provided information for this synthesis, often after discussions with local stakeholders over their experiences.

### *0.16 Cormorant population modelling*

The most well-supported Cormorant population model scenarios using current information indicated three important things. First, that the effect of culls at the 1998-9 level (i.e. 17, 000 birds shot) was limited. Second, that increasing the annual cull to 30, 000 birds would have limited effect at the population level. Third, that shooting 50, 000 birds per year was predicted to lead to population extinction in 20-40 years. The modelling approach also demonstrated that increasing the number of culled Cormorants was risky because once the compensatory power of the population is overcome, it will inevitably decline towards extinction if the cull is unchecked. One

general inference was that culls should be planned so that they become the most powerful density-dependent mechanism affecting the target population. This strategy would require a well parameterised population model and should also be accompanied by monitoring programmes. Even though Cormorant population control through culling is feasible it may not be the most efficient, economical or ethical way of limiting Cormorant damage to fisheries, and other interests, across Europe. Research suggests several limitations to culling and these are discussed.

#### *0.17 Relatively large-scale Cormorant control*

The synthesis of general information on actions against Cormorants included information from all 25 countries covered by the REDCAFE project. Some form of national or regional Cormorant management plan was in effect in 11 of these countries. A further four countries had a legal regulation in effect that allowed Cormorant culling. Overall, such a regulation was in effect in 14 countries. In a further 6 countries licences could be obtained for the limited killing of Cormorants at particular sites as a aid to scaring. In most countries (84%), there was either no killing of Cormorants or it was uncoordinated. Few countries (16%) had a co-ordinated culling programme. Few countries (or regions therein) provided either financial compensation for fish losses caused by Cormorants or financial aid for Cormorant exclosures or scaring programmes (16% and 24%, respectively). Of the 25 countries, ten recorded the destruction or disturbance of Cormorant colonies in recent (i.e. 1990-2002) years, with 102 colonies reported to be affected annually. As a result a minimum of 5,194 Cormorant nests were reported to be destroyed annually in five countries. Between 600-650 Cormorant nestlings were also reported to be killed in three countries. Numbers of both nests and nestlings destroyed were known to be under-recorded. Around 10, 000 adult Cormorants (of the 'Atlantic' *carbo* race) are hunted legally as game in Norway outside the breeding season. During this time of year, a further 18 countries reported killing Cormorants (mostly the 'Continental' *sinensis* race) as a control measure. Here, between 41-43, 000 adult birds (including young birds in their first winter) were reported to be killed annually. However, given the unprecedented number of Cormorants killed in France in 2001/02, and the fact that many of the birds killed were juveniles in their first winter, it is more appropriate to say that between 41-43, 000 fully grown birds were killed in 2001/02. A further 4,598 Cormorants were reported to be killed annually during the breeding season in six countries. However, this was known to be an underestimate. Over 248 night roosts were reported to be destroyed or damaged annually in nine countries. This figure was a considerable underestimate because roosts were also known to have been destroyed or disturbed in three other countries.

#### *0.18 Site-specific actions: non aquaculture habitats*

A total of 33 site-specific techniques used regularly to reduce the effects of Cormorants at feeding sites were recorded for 16 countries. However, only three techniques were used regularly at all five feeding habitats (small rivers, large rivers, small stillwaters, very large waterbodies, aquaculture): the use of live ammunition to scare birds, shooting birds to reinforce other forms of scaring, and shooting birds to reduce their numbers at specific sites. Eleven techniques were recorded in regular use on small and large rivers. Only two of these appeared to be effective in the long-term (i.e. years), although both of them (improving fish habitat quality and submerged fish refuges) were primarily related to the management of fishes rather than to that of Cormorants. Several other techniques appeared to be effective on rivers for months.

Eight techniques were recorded in regular use on small lakes. All appeared to be effective only for days, the exceptions being the use of two audio techniques (pyrotechnics/fireworks and live ammunition) and two lethal techniques (shooting to scare or to kill limited numbers of birds). Ten techniques were recorded in regular use on very large water bodies (lakes and coasts). Three audio techniques and three lethal Cormorant control techniques appeared effective over the time-scale of weeks to months. Other techniques appeared effective for only days.

#### *0.19 Site-specific actions: aquaculture habitats*

Twenty eight techniques were recorded in regular use at aquaculture facilities. Eight bird-proof barrier techniques appeared to be effective for up to years, although in some cases the same techniques were reported only to be effective for days. Alterations to fish stocking at aquaculture facilities appeared to be effective for up to months, as did the use of two audio techniques (pyrotechnics/fireworks and live ammunition) and three forms of lethal Cormorant control.

#### *0.20 Cormorant management tools: conclusions*

Very few techniques were, according to the experience in 16 countries covered by the synthesis, considered to be effective in the long-term (i.e. years). These long-term techniques appear to fall into two broad categories. First, those involving the alteration of fish habitat at some 'natural' rivers and lakes. Second, those involving the erection of various bird proof barriers (e.g. narrow mesh enclosures, wires, submerged anti-predator nets) at aquaculture facilities (both ponds and net pens/cages). Many other techniques used regularly can be effective for up to months at some sites. However, the same techniques were reported to be effective for only days, or not at all, at other sites. Overall, the practicability, acceptability and costs of all techniques used regularly were highly variable. The most likely explanation for such variation is that it is related to site-specific features. These are likely to be two-fold. First, the physical location of the site, its size, the type of fishery, the number of Cormorants involved etc. Second, the scale of the Cormorant 'problem' in financial terms.

#### *0.21 Cormorant-fishery conflict resolution: a case study*

REDCAFE analysed a specific Cormorant-fishery conflict case study, in the form of a three-day Workshop designed to give project participants and local stakeholders the opportunity to share their knowledge and experience. This case study also formed the basis for evaluating REDCAFE progress and the applicability of the 'REDCAFE experience' to the real world. Furthermore, it allowed participants to explore whether the project's concept of equitable stakeholder involvement was a useful framework for future Cormorant-fisheries conflict resolution elsewhere in Europe. An opportunity arose to link the project to a 'live' conflict case study - that of Cormorants and recreational fisheries in the Lea Valley, Hertfordshire, south-east England. Importantly, selecting the Lea Valley Cormorant-fishery issue also allowed REDCAFE to link with Fisheries Action Plans, and the government agency-led process being developed to address and prioritise issues affecting inland fisheries at a catchment scale. The REDCAFE case study was placed in perspective through reviews and discussions of values and dialogue in conflict resolution and management, Fisheries Action Plans in the UK, and the Lea Valley case study area.

### *0.22 Lea Valley Workshop*

Workshop delegates comprised 36 REDCAFE participants, representing 20 countries, and 16 stakeholders, representing 11 institutions or organisations. Successful conflict management depends on conflicting parties opening communication channels and developing networks of trust for effective collaboration and dialogue. REDCAFE thus worked closely during the Workshop with a facilitator skilled in environmental conflict management. The Workshop began the process of approaching the numerous environmental conflicts apparently affecting the Lea Valley. Although time was short, many important issues were addressed and developed, including conflict management experiences from both continental Europe and the Lea Valley itself. Several key issues arose from discussions with local stakeholders. First, many believe that the main problem facing the Lea Valley is an economic one. Economic measures of angling 'effort' (i.e. day and season ticket sales and angling club membership) have all fallen considerably in the last decade. This has had a knock-on effect on the local economy. Second, several lines of evidence suggest that many fish stocks and/or catches there have declined dramatically. The perception is that most small fish – both small individuals and small species - have declined, whilst there are still some fisheries containing large individuals (i.e. 'specimen' fish). There is also some evidence that the distribution of fish has changed within the Lea Valley. Third, the lack of fish, and the related economic decline, has local conservation implications, social implications, and planning and policy implications. These are all discussed.

### *0.23 Workshop progress*

Key local issues were summarised in an initial 'problem statement' for the Lea Valley. Substantial progress was made in identifying critical scientific and social issues in cormorant/fisheries conflicts. Cormorant-fishery conflicts play a part in the mix of issues facing the Lea Valley but one important outcome of the Workshop was to situate these conflicts in a broader social, economic and ecological context. Local stakeholders made considerable progress where escalating conflicts had become significant obstacles in the Lea Valley. REDCAFE participants had the opportunity to explore part of a conflict management process that related directly to many Cormorant-fishery conflicts across Europe. The Workshop process enabled significant progress to be made in several areas: (a) linking scientific processes and data to real-world social issues, (b) agreeing initial problem statements, stakeholders and needs, (c) identifying relevant agencies, people and pathways for action planning, and (d) identifying research priorities and dissemination actions that link the need for strong, evidence-based scientific knowledge with social and strategic planning needs.

### *0.24 Workshop evaluations*

A specific element of the REDCAFE project was to evaluate the conflict resolution Workshop in terms of determining whether the project's concept of equitable stakeholder involvement was a useful framework for future Cormorant-fisheries conflict resolution elsewhere in Europe. To this end, the Facilitator organised an anonymous questionnaire survey of delegates immediately after the Workshop. Twenty-six responses (50% of Workshop delegates) were received and almost all agreed that the case study was useful and enjoyable and that REDCAFE had helped them relate conflict management methods to Cormorant-fisheries conflicts elsewhere. A series of questions were also asked of delegates and those responding to the questionnaire provided over 200 responses which are synthesised in the report.

#### *0.25 The REDCAFE process: main strengths*

The most commonly cited strength of the case study Workshop, and of the REDCAFE process in general, was the development of trust between project participants and other stakeholders, and effective dialogue between scientists and others. Next followed the pan-European involvement and collaboration produced by the project and the opportunity it has provided to bring international perspectives to bear on local case studies. Another important strength identified was the project's attempts to reach consensus on Cormorant-fisheries conflicts through collaboration with social scientists.

#### *0.26 The REDCAFE process: main weaknesses*

In relation to the case study Workshop, the commonest weaknesses identified were lack of time and the involvement of too few local stakeholders. It was recognised that these constraints probably limited, to some degree, discussions on potential site-specific management tools. More generally, policy makers should have been included as REDCAFE participants and the continued need for effective dialogue between all interested parties was highlighted.

#### *0.27 The REDCAFE process: main lessons learned*

Several lessons for the REDCAFE project were recorded. The most frequent involved the vital importance of participation and dialogue. Almost all stakeholders stated that conflicts can only be resolved through relationships and trust: people must work together, ideally in face-to-face discussions, to develop solutions. All those involved in dialogue must consider the language they use and be aware that different participants (individuals or groups) will have different levels of confidence and enthusiasm. Respondents also noted that it takes time to understand conflict and decide how best to manage it. There may be no ultimate solutions but effective dialogue will invariably help to resolve conflicts. Another important lesson was that large-scale culling of Cormorants will almost certainly be ineffective. Cormorants are now an established element of many aquatic ecosystems and people need to learn to live with them. Scientific information is necessary to inform debate and potential mitigation policies, and REDCAFE has demonstrated that clear communication of scientific information can influence other stakeholders' perceptions and understanding and *vice versa*. Other important REDCAFE lessons were cited and these are discussed in detail in the report.

#### *0.28 Looking forward: overview*

REDCAFE has attempted to synthesise, for the first time, key stakeholder groups' views and perceptions on Cormorant conflicts with fisheries (and, to a lesser extent, with the wider environment) in a standardised way across Europe. Despite methodological limitations, many clear pictures emerged and these are discussed. Just as importantly, collecting and collating information for this synthesis has allowed REDCAFE participants (primarily natural scientists or those working closely with them) to forge links with local stakeholders experiencing conflict issues at first hand. REDCAFE offered the first opportunity to apply recognised conflict management techniques to Cormorant-fisheries interactions at the pan-European level. Through discussions with stakeholders it was clear that conflicts with Cormorants are not the only ones facing many fisheries and environmental stakeholders. To better understand the nature of Cormorant-fishery conflicts it is useful to consider other internal and external issues leading to conflicts over fisheries resources. These issues, both

environmental and social, are often complex and closely linked. Environmental conflicts over resources, including those involving fisheries, usually involve numerous issues. This appeared true across Europe: many of the stakeholders who provided specific information on Cormorant conflict issues for the present synthesis also described other issues, fears and concerns affecting their businesses or recreation. Many stakeholders also recorded concerns over the creation of sustainable fisheries and the development and implementation of effective, 'holistic' fisheries management programmes. Some of the other wider concerns affecting fishermen contributing to the present synthesis related to ownership and property rights and to changes in market economies. These issues are discussed in the report. The evaluation process confirmed that the REDCAFE philosophy of developing interdisciplinary links within and between the fields of natural and social science was very useful. Moreover, the project clearly demonstrates the necessity, and value, of dialogue and participation between all stakeholders (or their legitimate representatives) involved in Cormorant-fishery conflicts. Evaluations also showed that REDCAFE's approach to a specific Cormorant-fishery conflict case study provides a useful framework for similar activities elsewhere. There is acknowledgement that the process of conflict management will take time and require appropriate resources, including funds.

#### *0.29 Looking forward: case studies, individuals and stakeholder groups*

At the local level, by far the most commonly anticipated next step was to consider potential site-specific management techniques based on lessons learned from the REDCAFE synthesis. There is a strong desire to put theories into practice and to try mitigation measures that have been shown to work elsewhere. For many, next steps should include exploring the possibilities of developing and implementing local fishery management, or action, plans for specific case studies and/or the building of partnerships at the national level between fishery and conservation organisations such as the Moran Committee in the UK. REDCAFE emphasised the importance of making concerted efforts to create participation, dialogue and consensus building between local stakeholders involved in Cormorant-fisheries conflicts across Europe. This will require effective dissemination of relevant information at local, regional, national and international levels. Politicians and policy makers should also be included in such dissemination activities.

#### *0.30 Looking forward: the scientific community*

While social issues now feature strongly in the minds of the natural scientists involved in the REDCAFE project, many in that community expressed clear needs to further improve understanding of ecological issues. Scientists also realise the need to forge better links with others. Although scientific independence and rigour remain crucial, there is a need for scientists to apply their research results to real life cases. Scientists also need to collaborate with other stakeholders and local people, for example in the development of local management plans. Such collaboration will require scientists to communicate practical information to others in a clear manner and to maintain dialogue with all interested parties. Natural and social scientists also need to forge closer links because Cormorant-fisheries conflicts are situated in social and political contexts.

#### *0.31 Looking forward: Fisheries co-management*

While REDCAFE focused on Cormorant-fishery conflicts, other tensions were recognised by the project as influencing them. Addressing such broad fisheries

conflict issues is not trivial and will take time and require trust between stakeholders. Furthermore, in order to avoid inadequate fisheries policies and management systems, that tend to treat the symptoms rather than address underlying problems, broader environmental and institutional factors should be taken into account and fundamental socio-cultural conditions must also be given high consideration. Participatory co-management in fisheries, where managers and local fishermen co-operate in drafting policy, may facilitate successful management while also offering the possibility of reducing public costs. If natural resource management is to be sustainable in the long term, an understanding of human behaviour is vital and this multidisciplinary approach was recognised by REDCAFE. The fundamental challenge for fisheries management in this context is to find ways of expanding technical expertise whilst increasing collaboration in decision-making processes. In the past there has been much co-operation between fishermen and scientists at the individual level but a more organised management structure is required to bring these, and other, groups together. REDCAFE's work established an area of co-operation between natural scientists, local environmental stakeholders (fishermen and conservationists) and policy makers which should form the basis of future dialogue and collaboration.

### *0.32 Looking forward: future research*

A major challenge for natural scientists will be to make their work more relevant and useful to stakeholders. It is clear that different stakeholders involved in Cormorant-fisheries conflicts have different values and perceptions over these issues. It is also clear that other stakeholders view scientists as having different values and perceptions. Thus, scientists should be considered as another stakeholder group involved in the issue of Cormorants and fisheries. Given the recognition that there is no single value or perception (i.e. 'reality') for all the different stakeholders groups within this conflict, it is unrealistic to expect a single method of collecting, analysing and interpreting useful scientific information. The development of a rigorous scientific research programme to address Cormorant conflict issues will have to maintain high scientific standards but will also have to be both relevant to and influential in the decision-making process. There is a need for a practical pan-European Cormorant-fishery research programme that includes ecological study, collaboration between natural and social scientists and a strong conflict management element. Similarly, there is a need for long-term studies to quantify the effectiveness of various measures to mitigate against Cormorant problems at fisheries. Stakeholders have a long list of possible management actions against Cormorants but relatively little guidance on their likely effectiveness, practicability, acceptability or costs at a specific site. Clearly, considerably more work is required to trial the use of techniques to reduce Cormorant impact at feeding sites. Whatever framework future scientific research into Cormorant conflicts takes, it is clear that all stakeholders are concerned over the common issues of quality, health and status of biological resources in wetland systems. Dialogue with stakeholders highlighted several areas where major conflicts were currently poorly served by scientific literature and these are discussed. However, it must be stressed that such research should be undertaken with participation from stakeholders at all stages where possible. Ultimately, this should increase the useful knowledge of both scientists and other stakeholder groups whilst also increasing collaboration between all parties, but particularly local people, in the decision-making process with regard to Cormorant conflict issues across Europe.

### *0.33 Looking forward: concluding remarks*

Full information from REDCAFE should be disseminated as widely as possible so that the lessons learned from the project can be applied elsewhere. The establishment of a pan-European information exchange network would greatly facilitate the conflict resolution process and allow stakeholders to view their own particular situations in the broader continental context. Information must be exchanged at several levels: within and between disciplines of natural and social science, between scientists and other stakeholders, and between all interested parties and politicians, policy makers and the general public. The most important next step after dissemination is to build on the findings of REDCAFE so that local stakeholders can begin to develop effective site-specific strategies for resolving local conflicts. The formation of an information exchange network would be a very useful tool to facilitate the rapid transfer of ideas, experiences, management techniques, their implementation and subsequent outcomes. It could also offer stakeholders opportunities for discussion and could provide them with clear information on the actual costs (both invested and saved) of specific techniques. Although the REDCAFE project is the most comprehensive attempt to address Cormorant-fishery conflicts at the pan-European scale, it is clear that the project is merely the first step. Opportunities must now be explored to further develop the foundation framework that REDCAFE has developed in linking science with society and advancing processes of conflict management across a range of European contexts.

The REDCAFE Cormorant-conflict synthesis demonstrated clearly that such conflicts are complex, in terms of both biology and equally important social and economic issues. This synthesis is an important first stage towards developing trust and collaborations between all those affected by Cormorant conflicts. These issues are as much a matter of human interests as they are of biology. It is hoped that this element of REDCAFE's work will indeed be the start of a management process for Cormorant-fisheries conflict issues and, by implication, for wider environmental issues affecting fisheries and aquatic conservation across Europe. A formal approach to applying REDCAFE philosophy to the thousands of other case studies across Europe is needed. Moreover, the onus is currently on biologists to solve what are essentially people-people conflicts, professionals in other disciplines should be increasingly involved in these conflict management issues.

# 1 Introduction: background and aims of the project

## 1.1 Pan-European Great Cormorant populations

Two subspecies of Great Cormorant (hereafter ‘Cormorant’) occur in Europe: the ‘Atlantic’ subspecies *Phalacrocorax carbo carbo* and the ‘Continental subspecies *P. c. sinensis*. Latest (1995) breeding estimates for *carbo* are of 40,000 pairs, mostly on the coasts of Norway, UK, Ireland and northern France, representing over 80% of the world population of the nominate race (Debout *et al.* 1995). Although there are no estimates for *sinensis* populations during the 19<sup>th</sup> century or the first half of the 20<sup>th</sup>, it is likely that numbers in the remainder of Europe had declined to an unprecedented level of around 800 breeding pairs in the Netherlands in the early 1960s. Thereafter, numbers increased dramatically to over 150,000 pairs throughout the region in 1995 (van Eerden & Gregersen 1995) and it is likely that the species is now more numerous than ever before.

The geographical range of these populations has also expanded with Cormorants returning to some areas after a long absence whilst also moving into areas previously never occupied. Recent DNA studies have shown one consequence of such population increases and associated range expansion. *Sinensis* birds are breeding in inland colonies in the UK, living sympatrically, and probably hybridising, with *carbo* populations there (Goostrey *et al.* 1998).

The reasons for such expansion are unclear but possible causal factors include a “non-limiting food supply” (i.e. populations are not limited by a lack of food), protection of breeding sites and reduction in persecution throughout Europe (van Eerden & Gregersen 1995, Bregnballe & Gregersen 1997). The expansion of the European Cormorant population must be considered in the context of unprecedented landscape and social changes during the late 20<sup>th</sup> century. For instance, industrial activity in western Europe created many wetland habitats as a result of such things as gravel extraction and the construction of reservoirs. On the other hand, the recent decline in heavy industry in east/central Europe has led to reductions in aquatic pollution and an associated recovery of fish populations. Furthermore, the aquaculture industry has expanded across the whole continent, leading to areas of intensive freshwater fish production and enhanced stocks through the release of hatchery-reared fish.

Undoubtedly, protective legislation, particularly EEC Directive 79/409 on the Conservation of Wild Birds, has also been an extremely important factor in the increase of Cormorant populations throughout the region (van Eerden *et al.* 1995). Other important protective legislation in the EU includes the Bern Convention on the Conservation of European Wildlife and Natural Habitats and the Bonn Convention on the Conservation of Migratory Species of Wild Animals and, in the EU and elsewhere, the Ramsar Convention on Wetlands of International Importance especially as waterfowl habitat.

## 1.2 Basic conflicts between Cormorants and fisheries

Cormorants are generalist fish-eating predators taking a wide variety of species in shallow coastal seas, freshwater fisheries (natural and stocked artificially) in lakes and rivers, and both traditional/extensive and intensive aquaculture systems (Cramp & Simmons 1977). In almost all countries where Cormorants occur, their increasing

numbers and geographical spread has led to a growing number of conflicts with commercial fisheries and recreational angling interests (e.g. Bildsøe *et al.* 1998 and Suter 1995, respectively).

These conflicts arise either through direct consumption of commercial or rare fish species or through fears of indirect effects such as injury to fish and the spread of diseases and/or parasites that increase fish mortality and reduce their market value. There are clear cases of Cormorant damage to fishing gear and ensnared fish, as well as documented cases of considerable impact at fish farms and small water bodies (see van Dam & Asbirk 1997). Demonstrating the impact of Cormorants in large rivers and other water bodies is difficult because of ecological complexities. Nevertheless, annual losses as a result of Cormorant predation have been variously estimated at over 4 million ECU for European fishery yields in 1992 (Adamek *et al.* 1997) and at 163.7 million ECU for losses of commercial fish during the winter in relation to recreational angling, a sport for at least 23 million EU citizens (EAA, 1998).

### *1.3 International legislation, conservation and management plans*

Given these conflicts, where a species causes “*serious damage*” to specified interests such as fisheries and where other satisfactory solutions are lacking, several European Member States have derogated from their protective provisions with regard to the Cormorant under Article 9 of the EU Bird Directive. Although Article 9 derogations, or a national equivalent, have been applied on a local scale, the calls for further measures to reduce the population size of Cormorants made by some fisheries interests, particularly those in regions where Cormorants over-winter, increased during the 1990s. In particular, the governments of Denmark and the Netherlands - the countries where ca. 36% of the European Cormorants breed - were urged to address this issue. Thereafter, discussions were aimed at placing Cormorant management in the international legal framework of the Bonn Convention (Conservation of Migratory Species of Wild Animals). The underlying ethos was that the Cormorant, although not currently at risk, might be threatened again by illegal killings, if not properly managed. Moreover, there was also a need to protect two endangered cormorant species in the region, the Pygmy cormorant *P. pygmaeus* and the Socotra cormorant *P. nigrogularis*. These discussions resulted in a draft recommendation on the management of cormorants in the African-Eurasian region being presented to, and adopted by, the Conference of Parties at the fourth meeting of the Bonn Convention in 1994 (see van Dam & Asbirk 1997).

In 1996, the European Parliament adopted a ‘resolution on the cormorant problem in European fisheries’ (see van Dam & Asbirk 1997) considering it appropriate to take special temporary measures by means of scientific projects approved by the European Commission. These measures should aim at reducing the Cormorant’s impact on the environment by, for example, preventative action to restrict the reproduction of Cormorants and by the temporary exclusion of *P. c. sinensis* from Annex 1 of the Bird Directive. This resolution also called on the Council to take effective action to restore depleted fish stocks and ensure that the Common Fisheries Policy maintains fish at levels that can support both human fisheries and natural predators.

Under the auspices of the Bonn Convention, there followed a workshop ‘*Towards an International Conservation and Management Plan for the Great*

*Cormorant*' (Lelystad, the Netherlands, October 1996). An international meeting of experts to complete an Action Plan for the Management of the Great Cormorant in the African-Eurasian Region (Copenhagen, Denmark, September 1997) was held in the following year. This Action Plan aimed to minimise the conflict between fisheries interests and the Cormorant by "*ensuring that best practice is followed in mitigating, preventing and reducing their reported impacts on fisheries, while maintaining a favourable conservation status for the species.*" The Action Plan also stated that Range States should try to achieve this, in the following order of preference, through (a) appropriate site-specific management, (b) local management and control of Cormorants, and (c) co-ordinated management and control of Cormorants between Range States.

The *Action Plan for the Management of the Great Cormorant in the African-Eurasian Region* was sent to all European Range States with a request to implement the recommendations included therein. Any response to the action Plan was left to the discretion of individual Range States. Individual Range States appeared to largely ignore the Action Plan and continued with their own regional or national cormorant mitigation policies based on national and international legislation. There was little evidence that advice was being made available to Range States on the implementation of the Action Plan, nor was there facilitation to co-ordinate its implementation at the international level. Thus, in practice, and certainly for many affected by the 'cormorant problem', the Action Plan appeared to be ineffectual.

#### *1.4 Aims of the project*

The REDCAFE project was designed to complement and develop the previous work described above. It also addresses several of the main uncertainties highlighted during the development of the Action Plan (see van Dam & Asbirk 1997). For example, (a) the total size, long-term trends and movements/dispersal of the European Cormorant population, (b) the lack of reliable estimates of the social/economic aspects of Cormorant conflicts with commercial and recreational fisheries, and (c) the lack of reliable estimates of the efficacy and cost-effectiveness of methods of Cormorant control. Project aims were realised through synthesising available Cormorant/fisheries information, through identifying methods of reducing the current Europe-wide conflict between Cormorants and fisheries interests, and through collating expert evaluations of their practical use. The project took a novel approach to delivering solutions to these problems by, for the first time, bringing together avian, fisheries and social scientists and many relevant 'stakeholders'<sup>1</sup> to discuss and report on these issues in a rigorous, co-ordinated and equitable manner.

#### *1.5 Project set up*

During the years 1995 through 1999, it became increasingly apparent to many 'Cormorant researchers' that further progress towards Cormorant-fisheries conflict resolution required three important elements: first, a genuine pan-European approach to the problem; second, better integration of avian and fisheries science; third, a link

---

<sup>1</sup> The word 'stakeholders' is a difficult one: it means different things to different people and it is not easily translated into some languages. In the context of this report, the term 'stakeholders' is taken to mean (a) people who are affected (either positively or negatively) by a particular problem or activity or (b) people who can influence (either positively or negatively) the outcome or end result of a particular process. For further details see (Ramírez 1999).

between biological and social scientists and constructive dialogue between these groups and other stakeholders. Realising this, an initiative was taken by a group of natural scientists to propose an international network in order to synthesise relevant knowledge and to further progress pan-European Cormorant-fisheries conflict resolution.

This proposal was for a Concerted Action under the EU's 'Quality of Life and management of Living Resources' Fifth Framework Programme. Concerted Actions are "*Designed to encourage collaboration between teams of interested researchers and other actors, combining their accumulated expertise in a research network, to find solutions to problems common to all European Member States, but with better chances of solution through European collaboration.*" and "*... can be considered when pooling of data would facilitate common interpretation of facts and contribute to the development of harmonised standards, procedures and methodologies...*"

The European Commission agreed to award a financial contribution of 100% of the costs for this proposal ('REDCAFE'), paying for co-ordination staff time and the costs of four international Workshops. REDCAFE ran for 24 months (01.12.00 - 30.11.02) and in the first phase of this project, available information on Cormorant conflicts with fisheries was synthesised. In the second phase, available information on Cormorant ecology, focussing on those factors leading to conflicts with fisheries, was synthesised. In the third phase, a set of potential management tools, from continental to site-specific, was compiled. In each of these phases, as well as synthesising available knowledge, critical deficiencies and uncertainties were also highlighted. In the fourth phase a specific Cormorant-fishery case study was selected as a model for conflict resolution and to determine whether a useful framework could be established for future conflict resolution elsewhere. Finally, the report was compiled, including detailed findings from each of the four phases of the project and relevant national and international overviews.

This report contains two volumes. Volume 1 (this volume) provides the REDCAFE framework, a regional, national and international synthesis of Cormorant-fisheries conflicts (Chapter 3), a synthesis of relevant Cormorant ecology (Chapter 4) and a synthesis of potential management tools in all, or most, countries (Chapter 5). It also provides background, description, analysis and evaluation of a 'real world' case study exercise involving conflicts between Cormorants and recreational angling interests in a catchment in south-east England (Chapter 6). Volume 2 includes a Chapter for each participating country, in which the local situation is described in as much detail as possible. Each of the volumes, and each individual country text, can be read as stand alone texts.

Major contributions to this report were drafted by individual participants of this Concerted Action, namely: David Carss (all Chapters), Erik Petersson (Chapter 3), Mariella Marzano (Chapters 3, 6, 7), Mennobart van Eerden (Chapter 4), Stef van Rijn (Chapter 4), (Harold Claassen Chapter 4), Thomas Keller (Chapter 5), Morten Frederiksen (Chapter 5) and Scott Jones (Chapter 6, 7) and Participants contributing information to the country reports in Volume 2 are indicated at the beginning of each report in that Volume. To all colleagues and friends participating in this project, the Editor extends grateful and heartfelt thanks for their enthusiasm and hard work. Additionally the Editor would like to thank the following people for their input to

REDCAFE Work Packages: Joep de Leeuw, Maarten Platteeuw, Terry Mansbridge, Dennis Meadhurst and Adrian Taylor. Thanks also to: Denise Wright and Malcolm Collie (CEH Banchory) for invaluable IT help and advice throughout the project; Ian Cowx (University of Hull International Fisheries Institute) for hospitality and invaluable help during the project's first Work Package meeting; administrative staff at CEH Banchory, RIZA (Lelystad) and NERI (Kalø/Horsens) for all their help; local stakeholders – fishermen, fisherwomen and conservationists – for making our field visits so rewarding.


## 2 Introduction: the REDCAFE project

The purpose of the REDCAFE project (December 2000 – November 2002) is to synthesise current Cormorant/fisheries information and to identify, describe and evaluate methods of reducing the current Europe-wide conflict between conservationists and fisheries interests. Each REDCAFE participant was invited to contribute as much as possible to the synthesis process and, through collaboration, to develop an effective working partnership between groups, some of whom are, or have been in the recent past, sceptical of each other. The project also includes analysis of a specific Cormorant-fishery conflict case study, giving project participants the opportunity to share their extensive knowledge and experience with local stakeholders and *vice versa*. With these aims in mind, a pan-European network of project participants was established comprising 49 participants representing 43 organisations from 25 countries (Figure 2.1) and included seven main stakeholder groups: commercial fishermen, recreational fishermen, aquaculturists, avian/wetland conservationists, fisheries scientists, avian ecologists and social scientists.



**Figure 2.1** The 25 countries participating in the REDCAFE project 2000-2002.

Like most conflicts surrounding natural resource management, Cormorant/fishery conflicts are very often conflicts between people wishing to conserve natural resources and people wanting to make a living from them. However, there is common ground: a healthy environment is vital for all. Scientists, policy makers and users, beneficiaries and the public, all ‘stakeholders’ (see footnote #1) in the environment, need to work together to ensure its sustainability. Only by developing such partnerships can research provide solutions. The success of the REDCAFE project depended on dialogue between all interested parties, as part of a stepwise process of conflict resolution/management (Figure 2.2). This model is just one of many tools available for developing collaborations in natural resource management (Buckles & Rusnak 1999). However, it was considered appropriate to this project because REDCAFE’s aims were to synthesise current Cormorant-fisheries knowledge, create dialogue between scientists and stakeholders, and to identify current gaps in understanding, rather than to embark on new areas of relevant research.



**Figure 2.2** The REDCAFE concept: steps to resolving ecological conflicts.

It should be recognised that, regardless of the nature of any available scientific ‘evidence’, the views or perceptions of some stakeholders (e.g. that predators compete with humans for commercial resources) often create and/or fuel natural resource conflicts. There are then four steps to resolving these conflicts. The first is for scientists and other stakeholders to work together to identify the problem. From this comes step two, for the researchers to provide data to improve understanding of the conflict. Thirdly, scientists and other stakeholders can evaluate the research together. This can lead to effective partnerships between all parties and, perhaps to additional research. The final step is to implement and monitor resolution measures or policies. Crucially, at every step, stakeholders must be aware that conflict resolution is well described as ‘harmonisation of the incompatible’.

It was also recognised that although some individuals and institutions had already progressed through several steps of this, or a similar, process, none had done so at more than a national level. Moreover, it was clear, in some countries at least, that many important stakeholder groups had lost confidence in natural scientists to deliver a solution to the ‘cormorant problem’. Finally, given the continued heated debates across Europe about this problem, it was apparent that there were still dialogue difficulties, poor information exchange and few effective partnerships between scientists and stakeholders. This has resulted in little progress toward conflict resolution at the pan-European level. Against this background and following the process outlined above, REDCAFE sought to bring as many relevant people together as possible to discuss Cormorant-fisheries conflicts in an equitable manner. This last point was important, as the project would only succeed if all stakeholders had a voice and the opportunity to state their case. Through such active participation, all REDCAFE participants - even those who may have felt disenfranchised at first - were able to begin the critical process of building trust and rapport that underpins successful natural resource conflict management.

The REDCAFE project addressed four main topics. In order to offer participants common ownership of the work of the project, each topic was discussed initially at an international meeting. Here, participants shared their expertise and experiences and agreed on the most appropriate methodology for each piece of work.

The four main REDCAFE topics were:

(1) **A Conflict Workshop** to: (1) synthesise available information on Cormorant conflicts with fisheries, in relation to (a) specific fishery types, (b) geographical and temporal trends, and (c) the commercial fish species involved, and (2) quantify, with uncertainties, the economic losses to fisheries as a result of Cormorant activity.

(2) **A Cormorant Ecology Working Group** meeting to synthesise available information on Cormorant ecology, focussing on those factors that lead to conflicts with fisheries.

(3) **A Management Tools Working Group** meeting to (1) identify and evaluate existing Cormorant population models with particular emphasis on (a) predicting the ultimate size and geographical distribution of the European Cormorant population and (b) determining the levels of control that would be necessary to reduce the overall

population size, (2) identify and evaluate potential site-specific management techniques, incorporating current information on their efficacy and cost-effectiveness.

(4) **A Conflict Resolution Workshop** to focus on a specific Cormorant-fishery conflict involving recreational angling in the Lea Valley, Hertfordshire, south-east England. Here, REDCAFE participants met with a number of local stakeholders and both groups shared expertise and knowledge. This case study formed the basis for evaluating REDCAFE progress and the ability of the REDCAFE experience to be applied to the real world. Furthermore, it allowed participants to explore whether the project's concept of equitable stakeholder involvement was a useful framework for future Cormorant-fisheries conflict resolution elsewhere in Europe.